arXiv:1208.1121v1 [cond-mat.stat-mech] 6 Aug 2012

Gaussian Equilibration

Lorenzo Campos Venuti and Paolo Zanardi

Department of Physics and Astronomy and Center for Quantum Information Science & Technology, University of Southern California, Los Angeles, California 90089-0484, USA

A finite quantum system evolving unitarily equilibrates in a probabilistic fashion. In the general many-body setting the time-fluctuations of an observable $\mathcal{A}$ are typically exponentially small in the system size. We consider here quasi-free Fermi systems where the Hamiltonian and observables are quadratic in the Fermi operators. We first prove a novel bound on the temporal fluctuations and then map the equilibration dynamics to a generalized classical XY model in the infinite temperature limit. Using this insight we conjecture that, in most cases, a central limit theorem can be formulated leading to what we call *Gaussian equilibration*: observables display a Gaussian distribution with relative error $\Delta\mathcal{A}/\overline{\mathcal{A}} = O(L^{-1/2})$ where L is the dimension of the single particle space. We prove the conjecture analytically for the magnetization in the quantum XY model, and numerically for a class of observables in a tight-binding model.

PACS numbers: 03.65.Yz, 05.30.-d

Out of equilibrium and equilibration dynamics of closed quantum systems have been recently at the center of a renewed and intense interest [1–7]. In particular the issue whether quantum integrability plays a key role in equilibration and, if so, which one, has been investigated by several authors [8–10]. In this paper we will address this problem along the lines of the probabilistic approach to quantum equilibration advocated in [6, 7, 11]. Here the central object is the (infinite time) full time statistics of the expectation value of a quantum observable. We will focus on a particular, yet very important class of quantum integrable systems: quasi-free fermionic systems. Using fairly general central-limit type arguments as well as explicit analytic examples we will argue that a sharp distinctive feature of these systems, as opposed to the general interacting ones, is an exponential enhancement of the amplitude of the temporal fluctuations of a quadratic observable around its mean value. This appears to be a precise and quantitative way to make sense of the common folklore that integrability leads to a poorer (or no) equilibration.

The system is initialized in the generic state ρ_0 with N particles and both the evolution Hamiltonian H and the observable A are quadratic in the fermionic operators [20]. The Hamiltonian is $H = \sum_{x,y} c_x^\dagger M_{x,y} c_y = c^\dagger M c$ (notation $c^\dagger = \left(c_1^\dagger, \ldots, c_L^\dagger\right)$, L number of sites). The general quadratic observable has the form $A = \sum_{x,y} c_x^\dagger a_{x,y} c_y = c^\dagger a c$. We will assume that $\|a\|_\infty = O(1)$ [21] as this guarantees that the expectation values of A scales at most extensively with the system size [22] The main object of investigation is $\mathcal{A}(t) = \mathrm{tr}\left(A e^{-itH} \rho_0 e^{itH}\right)$. Exploiting the Gaussian nature of the problem and introducing the covariance matrix $R_{y,x} := \mathrm{tr}\left(\rho_0 c_x^\dagger c_y\right)$ $(0 \le R \le \mathbb{1})$ one can show that the expectation value $\mathcal{A}(t)$ reduces to a trace in the one-particle space:

$$\mathcal{A}(t) = \mathrm{tr}\left(a e^{-itM} R e^{itM}\right) . \qquad (1)$$

Eq. (1) is perfectly analogous to its many-body version $\mathcal{A}(t) = \mathrm{tr}\left(A e^{-itH} \rho_0 e^{itH}\right)$ with R playing the role of the initial state ρ_0. There is however one importance difference: while $\mathrm{tr}\rho_0 = 1$ one has $\mathrm{tr}R = N = \nu L$, i.e. is *extensive* (we defined $\nu = N/L$ the filling factor).

For unitary evolution, the density matrix $\rho(t) = e^{-itH}\rho_0 e^{itH}$ does not converge neither in the strong nor in the weak topology [6]. Equilibration must be formulated in probabilistic terms. Given the observation time window $[0, T]$, the observable $\mathcal{A}(t)$ has probability $P_A(\alpha)\, d\alpha$ of being in the interval $[\alpha, \alpha + d\alpha]$, where the probability density is given by $P_A(\alpha) = \overline{\delta(\mathcal{A}(t) - \alpha)}^T$, and the time average operation is $\overline{f}^T = T^{-1}\int_0^T f(t)\, dt$. For simplicity we will always take the limit $T \to \infty$ when taking time average and write simply $\overline{f}$ in place of $\overline{f}^\infty$. Roughly speaking an observable A *equilibrates* if its probability density $P_A(\alpha)$ is highly peaked around its mean $\overline{\mathcal{A}(t)} = \mathrm{tr}\left(A\overline{\rho}\right)$. [6]. The role of equilibrium state is played by the time-averaged density matrix $\overline{\rho}$.

The question we are going to address here is: what is the size of the fluctuations of $P_A(\alpha)$ for observables A in this quasi-free setting?

Let us start with a few remarks on the nature of the equilibrium state itself. If the spectrum is non-degenerate, the average, dephased, state $\overline{\rho}$ has the form $\overline{\rho} = \sum_n p_n |n\rangle\langle n|$ where $|n\rangle$ are many-body eigenstate of H corresponding to energy E_n and $p_n = \langle n|\rho_0|n\rangle$. The powers of the Hamiltonian H^n, $n = 0, 1, \ldots, d-1$ are linearly independent if there are d different eigenvalues E_n and if the spectrum is non-degenerate d coincides with the Hilbert space dimension [23]. In this latter case the average state can always be written in the form $\overline{\rho} = \exp\left[\sum_{k=0}^{d-1} \alpha_k H^k\right]$, the so called generalized Gibbs ensemble [8]. The coefficients α_n depend on the initial state ρ_0 and on the eigenvectors $\{|n\rangle\}$. The condition to write the coefficients α_n in terms of the p_n is precisely that of the invertibility of the Vandermonde matrix $V_{n,k} = (E_n)^{k-1}$ $(n, k = 1, \ldots, d)$, i.e., once again, non-degeneracy of the spectrum, since $\det V = \prod_{n<k}(E_k - E_n)$. The relation expressing the p_n's in terms of the α_k's is: $p_n = \exp\left[\sum_{k=0}^{d-1} \alpha_k E_n^k\right]$. Since

$\overline{\rho} = \exp\sum_n \ln p_n |n\rangle\langle n| = \exp\sum_{k=0}^{d-1}\sum_n \alpha_k E_n^k |n\rangle\langle n|$, the inverse relation is, in vector notation $\boldsymbol{\alpha} = V^{-1}\ln \boldsymbol{p}$. The inverse of the Vandermonde matrix can be found explicitly by expanding the identity $|n\rangle\langle n| = \prod_{k\neq n}\frac{H-E_k}{E_n-E_k}$, multiplying by $\ln p_n$, summing over n and exponentiating see [24]

Let us now go back to the context of Eq. (1). The time averaged covariance $\overline{R} = \overline{e^{-itM}Re^{itM}}$ defines a Gaussian state $\rho_{\overline{R}}$, in fact $0 \leq R(t) \leq \mathbb{1}$ implies $0 \leq \overline{R} \leq \mathbb{1}$. Moreover, since $\mathrm{tr}R(t) = N$, for all t, $\mathrm{tr}\overline{R} = N$, $\rho_{\overline{R}}$ is a Gaussian state with N particles. Now, for what concerns quadratic observables of the kind $A = c^\dagger a c$, their time average expectation value is the same as that obtained with $\rho_{\overline{R}}$: $\overline{\mathcal{A}} = \mathrm{tr}a\overline{R} = \mathrm{tr}A\rho_{\overline{R}}$. In other words the states $\overline{\rho}$ and $\rho_{\overline{R}}$ are the same when restricted to quadratic observables. A generic Gaussian state can be written in the form $\rho_R = \mathcal{N}\exp\left(c^\dagger G(R) c\right)$ where $G(R) := \log[R^T(1-R^T)^{-1}]$ [12] and $\mathcal{N}$ is a normalization constant. From this it immediately follows that that $\rho_{\overline{R}}$ can be written as $\rho_{\overline{R}} = \mathcal{N}\exp\sum_k \lambda_k c_k^\dagger c_k$ where c_k's are eigenmodes of H and coincides with ρ_{FC} of [8]. When the system equilibrates i.e., $\mathcal{A}(t) \to \mathrm{tr}\left(A\rho_{\overline{R}}\right)$ this remark shows the validity of the GGE (in the sense of [8]) for *any* initial state ρ_0 and all quadratic observables.

A bound on the variance Assuming the non-resonant condition ($E_n - E_m = E_p - E_q$ implies $n = m$ and $p = q$ or $n = p$ and $m = q$), Reimann has shown [13] that the temporal fluctuations $\Delta\mathcal{A}^2 = \overline{\left(\mathcal{A}(t) - \overline{\mathcal{A}}\right)^2}$ satisfy $\Delta\mathcal{A}^2 \leq \mathrm{diam}(A)^2\,\mathrm{tr}\overline{\rho}^2$ where $\mathrm{diam}(A)$ is the maximum minus the minimum eigenvalue of A. Now *typically*, for most initial states ρ_0, the purity $\mathrm{tr}\overline{\rho}^2$ is exponentially small in the system size implying exponentially small fluctuations [5].

In the quasi-free setting the non-resonant condition does not hold. Let us then seek for the analogous of the bound of Reimann in our quasi-free case. Let the one-particle Hamiltonian have the following diagonal form $M = \sum_k \Lambda_k |k\rangle\langle k|$. The time averaged covariance matrix is then $\overline{R} = \sum_k \langle k|R|k\rangle|k\rangle\langle k|$. We also define $F_{k,q} = \langle k|a|q\rangle\langle q|R|k\rangle$. Assuming the non-resonance of the gaps of the one-particle spectrum, one gets $\Delta\mathcal{A}^2 = \mathrm{tr}F^2 - \sum_k (F_{k,k})^2 \leq \mathrm{tr}F^2 = \sum_{k,q}|\langle k|a|q\rangle|^2|\langle q|R|k\rangle|^2$. Now R is a non-negative operator and so induces a (possibly degenerate) scalar product which satisfies Cauchy-Schwarz inequality: $|\langle q|R|k\rangle|^2 = |\langle q|k\rangle_R|^2 \leq \langle q|q\rangle_R\langle k|k\rangle_R = \langle q|R|q\rangle\langle k|R|k\rangle$. This leads us to

$$\Delta\mathcal{A}^2 \leq \mathrm{tr}\left(a\overline{R}a\overline{R}\right) \leq \|a\|_\infty^2\,\mathrm{tr}\overline{R}^2 \tag{2}$$

Now, since $0 \leq \overline{R} \leq \mathbb{1}$, $\mathrm{tr}\overline{R}^2 \leq \mathrm{tr}\overline{R} = \mathrm{tr}R = N$, we finally obtain $\Delta\mathcal{A}^2 \leq \|a\|_\infty^2\,\nu L$. While equation (2) is the quasi-free analog of the Reimann's bound it implies some important differences with respect to the general (non-free) case. Consider the situation where the observable A is extensive. In the non-free case the diameter of A is extensive, i.e. $\mathrm{diam}(A) = O\left(L^D\right)$ in D spatial dimensions. Moreover, the minimum value of the purity $\mathrm{tr}\overline{\rho}^2$ is $1/d$ and so is exponentially small in the system size. In the quasi-free setting, instead, the minimum value of $\mathrm{tr}\overline{R}^2$ in Eq. (2) is $\min\mathrm{tr}\overline{R}^2 = N^2/L = \nu^2 L$, since $\overline{R}/N$ defines a density matrix for which the minimum purity is the inverse of the matrix dimension L. All in all, recalling that $\|a\|_\infty = O(1)$, for extensive observables, the minimum of the bound to the variance is $O\left(L^2 e^{-\gamma L}\right)$, $\gamma > 0$, (for systems of linear size L), whereas in the quasi-free setting one has $O(L)$. This seems to hint at the fact that fluctuations in the quasi-free setting are proportional to the system size and are hence much larger than in the non-free case where they are exponentially small in the volume. Of course Eq. (2) is just an upper bound, and nothing prevents, in principle, from having a much smaller variance. In the following, however, we will argue that the extensive behavior of the fluctuations, in quasi-free systems, is in fact quite general.

Mapping to a classical XY model Let us write again the generic expectation value (1) in the basis which diagonalizes M: $\mathcal{A}(t) = \overline{\mathcal{A}} + 2\sum_{k<q}|F_{k,q}|\cos\left(t\left(\Lambda_k - \Lambda_q\right) + \phi_{k,q}\right)$ with $\phi_{k,q} = \arg F_{k,q}$. To obtain information on the probability density $P_A(\alpha)$ we consider the generating function $\chi_A(\lambda) := \overline{e^{\lambda\left(\mathcal{A}(t) - \overline{\mathcal{A}}\right)}}$. Now we observe that *if the (one-particle) energies are rationally independent (RI)*, as a consequence of a theorem on the averages, the infinite time average of $\mathcal{A}(t)$ is the same as the uniform average over the torus $\mathbb{T}^L$. In this case the generating function $\chi_A(\lambda)$ is exactly given by the partition function of the generalized, classical XY model with energy $E(\boldsymbol{\vartheta}) = 2\sum_{k<q}|F_{k,q}|\cos\left(\vartheta_k - \vartheta_q + \phi_{k,q}\right)$ and inverse temperature $\beta = -\lambda$. The matrix $|F_{k,q}|$ defines the lattice of the interactions while the phases $\phi_{k,q}$ give the offset from which the angles are measured. Note that the behavior of the density $P_A(\alpha)$ is dictated by $\chi_A(\lambda)$ in a neighborhood of $\lambda = 0$ which corresponds to infinite temperature of the classical XY model.

It is not difficult to engineer a situation which exactly reproduces the standard XY model in D-dimension. For example, it suffices to consider the Hamiltonian $H = \sum_{\boldsymbol{x}}\mu_{\boldsymbol{x}}c_{\boldsymbol{x}}^\dagger c_{\boldsymbol{x}}$, with $\mu_{\boldsymbol{x}}$ RI ($\boldsymbol{x}$ is a point of a D-dimensional lattice), choose the observable $A = \sum_{\langle\boldsymbol{x},\boldsymbol{y}\rangle}c_{\boldsymbol{x}}^\dagger c_{\boldsymbol{y}}$ ($\langle\boldsymbol{x},\boldsymbol{y}\rangle$ indicates nearest neighbor) and initial (Gaussian) state $|\psi_0\rangle = L^{-1/2}\sum_{\boldsymbol{x}}|\boldsymbol{x}\rangle = c_{\boldsymbol{k}=0}^\dagger|0\rangle$. In this case the partition function $\mathcal{Z}$ (free-energy $\mathcal{F}$) of the classical D-dimensional XY model is precisely the characteristic function of the observable A: $\chi_A(\lambda) = \mathcal{Z}$ ($\ln\chi_A(\lambda) = \mathcal{F}$). In fact in this case $a_{\boldsymbol{x},\boldsymbol{y}} = \delta_{\langle\boldsymbol{x},\boldsymbol{y}\rangle}$ while $R_{\boldsymbol{x},\boldsymbol{y}} = 1/L$ so that $F_{\boldsymbol{x},\boldsymbol{y}} = L^{-1}\delta_{\langle\boldsymbol{x},\boldsymbol{y}\rangle}$ which defines the nearest neighbor hyper-cubic graph.

We would like to stress here that the one-particle space has a natural underlying geometric structure. For instance, the labels k, q represent points in momentum (real) space in a superfluid (localized) phase and the distance $|k-q|$ is well defined. Now, when the matrix elements $|F_{k,q}|$ decay sufficiently fast as $|k-q| \to \infty$ the corresponding XY model is well defined in the thermodynamic limit, i.e. the intensive free energy has a limit as $L \to \infty$. This happens for instance in case $|F_{k,q}|$ decays decays exponentially in $|k-q|$ or if one has $|F_{k,q}| \sim 1/|k-q|^\gamma$ with $\gamma > D$. When this is the case one has $\chi_A(\lambda) = \exp L^D \mathcal{F}(\lambda)$. Moreover one expects $\mathcal{F}(\lambda)$ to be analytic in the high temperature, $\lambda = 0$, limit, meaning that all the cumulants are extensive. From this we immediately draw the central limit theorem (CLT): as $L \to \infty$ the variable $(\mathcal{A}(t) - \overline{\mathcal{A}})/L^{D/2}$ tends in distribution to a Gaussian with zero mean and finite variance given by $\partial^2_{\lambda=0}\mathcal{F}(\lambda)$. We call this situation *Gaussian equilibration.* It is important to stress that one cannot have Gaussian equilibration in the non-free setting or all the cumulants would scale in the same way. Instead in the non-free setting one expects extensive average but variance exponentially small in the volume.

We will now further corroborate these arguments with two examples where indeed Gaussian equilibration can be proven or shown.

Quench on the quantum XY model The Hamiltonian is given in terms of Pauli spin operators $\sigma_i^{x,y,x}$ (we use periodic boundary conditions)

$$H = -\sum_{i=1}^{L}\left[\left(\frac{1+\gamma}{2}\right)\sigma_i^x\sigma_{i+1}^x + \left(\frac{1-\gamma}{2}\right)\sigma_i^y\sigma_{i+1}^y + h\sigma_i^z\right] \tag{3}$$

In the quench scenario, the initial state is the ground state $|\psi_0\rangle$ of the Hamiltonian with certain values of the parameters γ_0, h_0. The parameters are then suddenly changed and $|\psi_0\rangle$ is evolved with the Hamiltonian with parameters γ_1, h_1.

The model in Eq. (3) has been long used as a testbed for the study of quantum phase transitions (taking place at $h = 1$ and zero temperature) in many body systems, and more recently in the realm of out-of equilibrium unitary dynamics. See [14, 15] and the more recent monumental [16] for more details and references. A Jordan-Wigner transformation brings Eq. (3) to a quadratic form in Fermi operators. Since σ_i^z in terms of Fermi operators is $\sigma_i^z = 2c_i^\dagger c_i - 1$, the transverse, total magnetization $M(t) = \sum_i\langle\sigma_i^z(t)\rangle$ is a quadratic observable. Its expectation value in the quench setting is given by [15]: $M(t) = 2\sum_{k>0}\cos\vartheta_k^{(1)}\cos(\delta\vartheta_k) + \sin\vartheta_k^{(1)}\sin(\delta\vartheta_k)\cos\left(t\Lambda_k^{(1)}\right)$ where $\tan\vartheta_k^{(i)} = -\gamma_i\sin k/(h_i + \cos k)$, $\delta\vartheta_k = \vartheta_k^{(1)} - \vartheta_k^{(0)}$ and $\Lambda_k^{(i)} = 2\sqrt{(\gamma_i\sin k)^2 + (h_i + \cos k)^2}$ are the one-particle energies. The quasi-momenta are quantized according to $k = \pi(2n+1)/L$, $n = 0, 1, \ldots, L/2 - 1$. At this point it seems quite natural to expect that the energies $\Lambda_k^{(1)}$ are rationally independent. For example one can show that the numbers $\cos(k_n)$ $k_n = \pi(2n+1)/L$, $n = 1, 2, \ldots, (L-1)/2$ are rationally independent for L prime [7]. Given the form of the dispersion we may expect that the requirement that L is prime may be lifted.

Assuming rational independence of the one-particle energies $\Lambda_k^{(1)}$, the corresponding classical XY model has energy $E(\boldsymbol{\vartheta}) = \sum_{k>0} W_k\cos(\vartheta_k)$ with $W_k = \sin\vartheta_k^{(1)}\sin(\delta\vartheta_k)$. Each classical spin '$k$' interacts with an external field along a fixed axis with strength W_k. The partition function factorizes, each integral over ϑ_k giving a Bessel function $I_0(\lambda W_k)$ and we obtain $\overline{e^{\lambda(M(t)-\overline{M})}} = \exp\sum_{k>0}\ln(I_0(\lambda W_k))$. Clearly $M(t) - \overline{M}$ is a sum of independent random variables, each with zero mean and variance $W_k^2/2$. Now we have the following

Theorem For any value of parameters $(\lambda_0, h_0) \neq (\lambda_1, h_1)$, the variable $(M - \overline{M})/\sqrt{L}$ as $L \to \infty$ tends in distribution to a Gaussian with zero mean and variance $(2\pi)^{-1}\int_0^\pi (W_k^2/2)\,dk$.

We can prove the theorem by showing that the Lyapunov condition is satisfied so that the central limit theorem follows from Lindeberg's theorem (see e.g. [17]). Following the notation of [17] we have $s_L^2 = \sum_k \sigma_k^2 = \sum_{k>0} W_k^2/2 \to L(4\pi)^{-1}\int_0^\pi W_k^2\,dk$. Then, with $\delta = 1$ $\overline{|X_k|^{2+\delta}} = 4/(3\pi)|W_k|^3$. The Lyapunov's condition with $\delta = 1$ is the vanishing of the following quantity as $L \to \infty$:

$$\frac{1}{s_L^3}\sum_{k>0}\overline{|X_k|^3} \overset{L\to\infty}{\longrightarrow} \frac{1}{\sqrt{L}}\frac{16}{3\sqrt{\pi}}\frac{\int_0^\pi |W_k|^3\,dk}{\left(\int_0^\pi |W_k|^2\,dk\right)^{3/2}}. \tag{4}$$

Indeed the RHS of Eq. (4) goes to zero as $L \to \infty$ since $|W_k| \leq 2$ for all k and $|W_k| \neq 0$ for almost any k for $(\lambda_0, h_0) \neq (\lambda_1, h_1)$.

Remark For small quench close to a critical point, the function $W_k \simeq \sin(\vartheta_k^{(1)})(\partial\vartheta_k/\partial x)\,dx$ (x is the quenched variable, $x = \lambda, h$), becomes highly peaked (for instance, the peak is around $k = \pi$ close to the Ising critical point at $h = 1$ and W_k/dx diverges as $1/k$). For finite L and sufficiently small quench, few terms W_k dominate and one can obtain (for finite L) a non-Gaussian distribution. Indeed, as discussed in detail in [11], this is the case in general: for small quenches close to a quantum critical point, observables become a sum of few independent random variables and the distribution acquires a universal double-peaked form.

Tight binding model We consider here the 1-D tight binding model $H = \sum_x\left(c_x^\dagger c_{x+1} + \text{h.c.}\right)$ with twisted boundary conditions $c_{L+1} = c_1 e^{i\theta L}$ as proposed in [18]. As quadratic observable take A with diagonal one-particle matrix: $A = \sum_{x=1}^{\ell} c_x^\dagger c_x$. The system is initialized setting all the N particles say to the left of the chain, i.e. the initial covariance matrix is $R = \operatorname{diag}(1, 1, \ldots, 1, 0, \ldots, 0)$ with N ones on the diagonal. The observable A is extensive for $\ell = \alpha L$, and the thermodynamic limit is given by $\alpha = \ell/L$, $\nu = N/L$ constant and $L \to \infty$. The time evolved observable reads $\mathcal{A}(t) = \sum_{k,q} g_N(k-q)\, g_\ell(q-k)\, e^{-it(\Lambda_k - \Lambda_q)}$ where the function $g_\ell(\xi)$ is given by $g_\ell(\xi) = L^{-1}\sum_{x=1}^{\ell} e^{-ix\xi}$. The matrix $F_{k,q}$ depends only on the difference $k - q$: $F_{k,q} = f(k-q) = g_N(k-q)\, g_\ell(q-k)$. The eigen-energies are given by $\Lambda_k = 2\cos(k+\theta)$ and the quasimomenta can be considered quantized as $k = 2\pi n/L$, $n = 0, 1, \ldots, L-1$. For $\theta = 0$ the energies are degenerate as $\Lambda_k = \Lambda_{\pi-k}$ but for most of the θ, $\pi - k$ does not belong to the Brillouin zone and the energies are non-degenerate. In this case the average is $\overline{\mathcal{A}} = \sum_k g_N(0)\, g_\ell(0) = L\nu\alpha$. Still the Λ_k cannot be RI as $0 = \operatorname{tr}M = \sum_k \Lambda_k$, however it is likely that there are few relations among the energies. For instance Tasaki showed [18] that for L odd and most of the $\theta \in (0, \pi/L)$, the one-particle energies satisfy the non-resonant condition. This implies that, for most θ, the variance is given by

$$\Delta\mathcal{A}^2 = \sum_{k,q} \left| g_N\left(k-q\right) g_\ell\left(q-k\right)\right|^2 - \sum_{k} \left[g_N\left(0\right) g_\ell\left(0\right)\right]^2 .$$

The double sum above can be evaluated going back to real space with result $L^{-2}\left[N\ell^2 - \ell\left(\ell^2-1\right)/3\right]$ (assuming $N \geq \ell$ otherwise swap N with ℓ), hence also the variance is extensive and one has $\Delta\mathcal{A}^2 = L\left(\nu\alpha^2 - \frac{\alpha^3}{3} - \nu^2\alpha^2\right) + O\left(L^{-1}\right)$.

On the other hand, using methods borrowed from statistical mechanics, we can get an approximate analytic expression for the whole cumulant generating function. We first assume rational independence of the energies Λ_k so that the problem is mapped to an classical XY model. A single (or a few) relation among the energies is a sort of boundary condition for the classical model and is not expected to change the leading, bulk, term. The energy of the classical model for the translation invariant case is $E\left(\boldsymbol{\vartheta}\right) = 2\sum_{d=1}^{(L-1)/2}\left|f\left(d\right)\right|\sum_{x=1}^{L}\cos\left(\left(\tilde{\vartheta}_{x+d}-\tilde{\vartheta}_x\right)+\phi_d\right)$ ($\phi_d = \arg f\left(d\right)$ and $\tilde{\vartheta}_x$ is a periodic extension of ϑ_x, i.e. $\tilde{\vartheta}_{x+nL} = \tilde{\vartheta}_x$). Now we note that $f\left(d\right)$ is highly peaked around $d = 0$, so we approximate the energy keeping only the nearest neighbor term, i.e. $E\left(\boldsymbol{\vartheta}\right) \approx 2\left|f\left(1\right)\right|\sum_{x=1}^{L}\cos\left(\left(\tilde{\vartheta}_{x+1}-\tilde{\vartheta}_x\right)+\phi_1\right)$. This is precisely a one-dimensional (classical) XY model with periodic boundary condition (and off-set ϕ_1) and can be solved via transfer matrix method [19]. The partition function becomes $\mathcal{Z} = \mathrm{tr}\mathcal{K}^L$ where the transfer matrix operator is $[\mathcal{K}h(\vartheta_x)](\vartheta_{x+1}) = \int_0^{2\pi} d\vartheta_x e^{\lambda|f(1)|\cos\left(\tilde{\vartheta}_{x+1}-\tilde{\vartheta}_x+\phi_1\right)} h(\vartheta_x)/(2\pi)$. The transfer matrix is non-hermitian because of ϕ_1. Using the identity $e^{K\cos(\alpha-\beta)} = \sum_{p\in\mathbb{Z}} I_p\left(K\right) e^{ip(\alpha-\beta)}$ where I_p are Bessel functions (they satisfy $I_0 > I_1 > I_2\cdots$ and $I_p\left(K\right) = I_{-p}\left(K\right)$) one sees that plane waves $e^{im\vartheta_x}$ are eigenfunction of $\mathcal{K}$ with eigenvalue $I_m\left(2\left|f\left(1\right)\right|\lambda\right)e^{-im\phi}$. The largest eigenvalue in modulus, with $m = 0$, gives immediately the free energy: $\mathcal{F} = \ln\mathcal{Z}/L = \ln\left(I_0(\lambda\left|f(1)\right|)\right)$. As expected, in this approximation, the cumulant generating function is extensive and analytic in $\lambda = 0$. One then has again CLT in its standard form: the variable $(\mathcal{A}\left(t\right) - \overline{\mathcal{A}})/\sqrt{L}$, tends in distribution to a Gaussian in the thermodynamic limit with variance $\partial^2_{\lambda=0}\mathcal{F} = \left|f(1)\right|^2/2$. The Gaussian prediction is clearly confirmed by a numerical experiment see fig. 1.

Conclusions In this paper we addressed the question of equilibration in quasi-free Fermi systems. The initial state is a general state while the evolution Hamiltonian as well as the observable are quadratic in the Fermi operators. While for general non-free systems the variance $\Delta\mathcal{A}^2$ is typically exponentially small in the volume, we find that in the quasi-free setting (for extensive observables) both the mean and the variance are proportional to the volume. This hints at the possibility of an underlying central limit theorem, a circumstance that we termed *Gaussian equilibration*. In this case the properly rescaled observable becomes Gaussian in the large L limit, and in general the relative error satisfies $\sqrt{\Delta\mathcal{A}^2}/\overline{\mathcal{A}} \sim 1/\sqrt{L}$.



Figure 1: Full distribution for the observable $A = \sum_{x=1}^{\ell} c_x^\dagger c_x$ around its mean for $L = 201$, $N = 100$, $\ell = 101$. The histogram is obtained sampling $\mathcal{A}\left(t\right)$ at 240,000 random times uniformly distributed in $[0, T_{\max}]$ with $T_{\max} = 180,000$. The thick curve is a Gaussian with zero mean and variance $\sigma^2 = 0.0208L$ as computed in the text.

We prove Gaussian equilibration for the magnetization in the quantum XY model and give evidence for a class of observables/initial states evolving with a tight-binding model. The key insight is a mapping of the equilibration dynamics to a generalized classical XY model at infinite temperature. As a by product we obtain a quantum setting (initial state, Hamiltonian, and observable A_{XY}), such that the equilibration dynamics of A_{XY} gives the solution of a classical XY model in D-dimension, and vice-versa.

Ultimately the origin of the exponential decrease of the signal to noise ratio illustrated in this paper lies in the exponential reduction of the effective phase space entailed by the quasi-free nature of our setup i.e., the many body space gets effectively replaced by the single particle one. A natural question for future research is whether a similar mechanism is at place for a more general class of integrable systems. For example it is tempting to conjecture that this might be indeed the case for systems integrable by Bethe Ansatz as interacting eigenstates are described in terms of single-particle quasi momenta (rapidities).

Finally, we would like to stress that the enhancement of the temporal fluctuations of $\Delta\mathcal{A}^2$ (for a quadratic observable A) should provide a universal and *experimentally testable* signature of integrability in the context of out of equilibrium dynamics.

The authors acknowledge partial support by the ARO MURI grant W911NF-11-1-0268. PZ also acknowledges partial support by NSF grants No. PHY-969969 and No. PHY-803304.

[1] T. Kinoshita, T. Wenger, and D. S. Weiss, Nature **440**, 900 (2006), ISSN 0028-0836, URL `http://www.nature.com/nature/journal/v440/n7086/abs/nature04693.html`.

[2] P. Calabrese and J. Cardy, Physical Review Letters **96**, 136801 (2006), URL `http://link.aps.org/doi/10.1103/PhysRevLett.96.136801`.

[3] S. Hofferberth, I. Lesanovsky, B. Fischer, T. Schumm, and J. Schmiedmayer, Nature **449**, 324 (2007), ISSN 0028-0836, URL `http://www.nature.com/nature/journal/v449/n7160/full/nature06149.html`.

[4] M. Rigol, V. Dunjko, and M. Olshanii, Nature **452**, 854 (2008), ISSN 0028-0836, URL `http://dx.doi.org/10.1038/nature06838`.

[5] N. Linden, S. Popescu, A. J. Short, and A. Winter, Physical Review E **79**, 061103 (2009), URL `http://link.aps.org/doi/10.1103/PhysRevE.79.061103`.

[6] L. Campos Venuti and P. Zanardi, Physical Review A **81**, 022113 (2010), URL `http://link.aps.org/doi/10.1103/PhysRevA.81.022113`.

[7] L. Campos Venuti, N. T. Jacobson, S. Santra, and P. Zanardi, Physical Review Letters **107**, 010403 (2011), URL `http://link.aps.org/doi/10.1103/PhysRevLett.107.010403`.

[8] M. Rigol, V. Dunjko, V. Yurovsky, and M. Olshanii, Physical Review Letters **98**, 050405 (2007), URL `http://link.aps.org/doi/10.1103/PhysRevLett.98.050405`.

[9] C. Kollath, A. M. Läuchli, and E. Altman, Physical Review Letters **98**, 180601 (2007), URL `http://link.aps.org/doi/10.1103/PhysRevLett.98.180601`.

[10] T. Barthel and U. Schollwöck, Physical Review Letters **100**, 100601 (2008), URL `http://link.aps.org/doi/10.1103/PhysRevLett.100.100601`.

[11] L. Campos Venuti and P. Zanardi, Physical Review A **81**, 032113 (2010), URL `http://link.aps.org/doi/10.1103/PhysRevA.81.032113`.

[12] I. Peschel, Journal of Physics A: Mathematical and General **36**, L205 (2003), ISSN 0305-4470, 1361-6447, URL `http://iopscience.iop.org/0305-4470/36/14/101`.

[13] P. Reimann, Physical Review Letters **101**, 190403 (2008), URL `http://link.aps.org/doi/10.1103/PhysRevLett.101.190403`.

[14] E. Lieb, T. Schultz, and D. Mattis, Annals of Physics **16**, 407 (1961), ISSN 0003-4916, URL `http://www.sciencedirect.com/science/article/pii/0003491661901154`.

[15] E. Barouch, B. M. McCoy, and M. Dresden, Physical Review A **2**, 1075 (1970), URL `http://link.aps.org/doi/10.1103/PhysRevA.2.1075`.

[16] P. Calabrese, F. H. L. Essler, and M. Fagotti, arXiv:1204.3911 (2012), URL `http://arxiv.org/abs/1204.3911`.

[17] P. Billingsley, *Probability and Measure* (John Wiley & Sons, 2012), ISBN 9781118341919.

[18] H. Tasaki, arXiv:1003.5424 (2010), URL `http://arxiv.org/abs/1003.5424`.

[19] D. C. Mattis, Physics Letters A **104**, 357 (1984), ISSN 0375-9601, URL `http://www.sciencedirect.com/science/article/pii/0375960184908168`.

[20] Throughout the paper we use interchangeably the terms quasi-free or quadratic, for observables quadratic in Fermi operators.

[21] For example, in the translation invariant case it suffices to have $a_{x,y} = a\,(x-y)$ sufficiently fast decaying.

[22] By diagonalizing a and exploiting unitary invariance of the operator norm, one finds $\langle A\rangle \leq \|A\| = \|\sum_\mu \alpha_\mu c_\mu^\dagger c_\mu\| \leq \sum_\mu |\alpha_\mu| \|c_\mu^\dagger c_\mu\| = \sum_\mu |\alpha_\mu| \leq \|a\|_\infty L$. Here the α_μ's are the eigenvalues of A and the c_μ's the fermionic operators associated to the corresponding eigenvectors.

[23] Note that one can have a non-degenerate spectrum even in free systems. It suffices, e.g., to have a rational independent one-particle spectrum.

[24] Explicitly $\sum_n p_n |n\rangle\langle n| = \exp\left[\sum_n \sum_{k=0}^{d-1} \frac{\gamma_{k,n}}{\Delta_n} \ln p_n\, H^k\right]$ where $\gamma_{k,n} = (-1)^{d-1-k} \sum_{j_1<\cdots<j_{d-k};j_i\neq n} E_{j_1} \cdots E_{j_{d-k}}$, and $\Delta_n = \prod_{j\neq n} (E_n - E_j)$.